## A Few Scientific Papers using Accutase/Accumax

Efficient Propagation of Single Cells Accutase-Dissociated human Embryonic Stem Cells, Bajpai, et al, <u>Journal of Molecular</u> <u>Reproduction and Development</u>, DOI 10.1002/mrd:1-10, 2007.

Human Embryonic Stem Cell-derived Dopaminergic Neurons Reverse Functional Deficit in Parkinsonian Rats, Yang, et al, <u>Stem Cells</u>,0: 2007-0494v1, 2007.

Efficient gene delivery and Silencing of mouse and human pancreatic islets, Lefebvre et al, BMC Biotechnology, 10:28, 2010.

Neural progenitors derived from human embryonic stem cells are targeted by allogeneic T and natural killer cells, Preynat-Seauve, et al, Journal of Cellular and Molecular Medicine, Vol. 13:9b, pp 3556-3569, March 2009.

In vitro cultivation and cryopreservation of duck embryonic hepatocytes, <u>Journal of Virological Methods</u>, Volume 157, Issue 1, pp 25-31, 2009.

Highly Efficient Differentiation and Enrichment of Spinal Motor Neurons Derived from Human and Monkey Embryonic <u>Stem</u> <u>Cells</u>, Tamaki Wada, et al, <u>PLoS One</u>, 4(8); e6722, 2009.

Human oligodendrocytes from embryonic stem cells: conserved SHH signaling networks and divergent FGF effects, Bao-Yang Hu et al <u>Development</u> 136, pp1443-1452, , 2009.

Restricted Spontaneous In Vitro Differentiation and Region-Specific Migration of Long-Term Expanded Fetal Human Neural Precursor Cells After Transplantation Into the Adult Rat Brain, Maciaczyk et al, <u>Stem Cells and Development</u>, Vol.18(7): 1043-1058, 2009.

Neural conversion of human embryonic stem cell colonies in the presence of fibroblast growth factor-2, Benzing, et al, <u>NeuroReport</u>, Vol 17, Issue 16, November 2006.

Human ES cell-derived neural rosettes reveal a functionally distinct early neural stem cell stage, Elkabetz, et al., <u>Genes &</u> <u>Development</u>, 22: 152-165, 2008,.

**ROCK inhibitor improves survival of cryopreserved serum/feeder-free single human embryonic stem cells,** Li, *et al.* <u>Human Reproduction</u>, Mar, 01; 24(3):580-9 2009. "Accutase significantly increases the survival of single hES cells after thawing."

Creation of Engineered Human Embryonic Stem Cell Lines Using phiC31 Integrase, Thyagarajan, et al, <u>Stem Cells</u>, Vol. 26 Issue 1, 119-126, 2007.

A Scaleable and Defined System for Generating Neural Stem Cells from Human Embryonic Stem Cells, Joannides, et al, <u>Stem Cells</u>, Volume 25, Issue 3, 731-737, 2006.

**Canine hemangiosarcoma originates from hematopoietic precursors with potential for endothelial differentiation**, Lamerota-Kozicki *et al.*, <u>Experimental Hematology</u>, Vol. 34 Pages 870-878, April 2006.

The JAK3 inhibitor WHI-P154 prevents PDGF-evoked process outgrowth in human neural precursor cells, Richards *et al.*, Journal of Neurochemistry, Vol. 97 Page 201, April 2006.

High Efficacy of Clonal Growth and Expansion of Adult Neural Stem Cells, Frank-Peter Wachs et al, Lab Investigation, 83:949–962, 2003.

Derivation of pluripotent epiblast stem cells from mammalian embryos, Gabrielle M. Brons et al, <u>Nature</u>, 448, 191-195 (12 July 2007).

Somatic Stem Cell Marker Prominin-1/CD133 Is Expressed in Embryonic Stem Cell-Derived Progenitors, Gabriela Kania, Stem Cells, Vol. 23 NO: 6, Pages: 791-804, 2005

Nuclear factor-KB controls the reaggregation of 3D neurosphere cultures in vitro, Widera et al., European Cells and Materials, Vol. 11, Pages 76-85, 2006.

Autologous adult rodent neural progenitor cell transplantation represents a feasible strategy to promote structural repair in the chronically injured spinal cord, Pfeifer, <u>Future Medicine</u>, Pages 255-266, July 2006.

Neural Conversion of Human Embryonic Stem Cell Colonies in the Presence of Fibroblast Grown Factor-2, Benzing *et al.,* <u>Developmental Neuroscience</u>, Vol. 17, No. 16, Pages 1675-1678, November 2006.

Lack of response to epidermal growth factor in adult neural progenitor cells, Kalluri *et al.*, <u>Molecular Neuroscience</u>, Vol. 16, Pages 835-838, May 2005.

Behavioral Changes in Unilaterally 6-Hydroxy-Dopamine Lesioned Rats After Transplantation of Differentiated Mouse Embryonic Stem Cells Without Morphological Integration, <u>Stem Cells</u>, Vol. 22 Pages 396-404, 2004.

High efficacy of clonal growth and expansion of adult neural stem cells, <u>Lab Investigation</u>, Vol. 83(7):949-962, . July 2003. "In addition, increased cell survival was obtained when Accutase, instead of trypsin, was used for enzymatic dissociation of NSC cultures."

**Oxygen Reduce Accumulation of type IV Collagen in endothelial Cell Subcellular Matrix via Oxidative Stress**, T. Brevig. et al, <u>Artificial Organs</u>, Volume 30 Issue 12, Pages 915-921, December 2006.

Circulating fibroblasts, Philips, Journal of Clinical Investigation, Vol. 114, Pages 438-446, August 2004.

Adult neural progenitor cell grafts survive after acute spinal cord injury and integrate along axonal pathways, <u>European</u> Journal of Neuroscience, Vol, 18 Issue 4, Page 743, August 2003.

Inhibition of death-receptor mediated apoptosis in human adipocytes by the IGF-1/IGF-1R autocrine circuit, <u>Endocrinology</u>. December 22, 2003.

Binding of gastrointestinal tumor cells to endothelial E- and P-selectin adhesion receptors leads to transient downregulation of sLeX ligands in vitro, International Journal of Colorectal Disease, Volume 18, Number 4, Pages: 292 – 299, July 2003.

A microcarrier cell culture process for propagating rabies virus in Vero cells grown in a stirred bioreactor under fully animal component free conditions. Rourou, *et al*, Vaccine, 10; 25(19):3879-89, 2007.

A cell-detachment solution can reduce background staining in the ELISPOT assay, Grant A, et al., <u>Methods of Molecular</u> <u>Biology</u>, 302:87-94, 2005.

**Oxygen Reduces Accumulation of Type IV Collagen in Endothelial Cell Subcellular Matrix via Oxidative Stress,** T. Brevig, et al, Artificial Organs, Volume 30 Issue 12 Page 915-921, December 2006.

Integrin Signalling Regulates the Nuclear Localization and Function of the Lysophosphatidic Acid Receptor-1 (LPA<sub>1</sub>) In Mammalian Cells, Waters *et al.* Biochemical Journal Immediate Publication, May 2006.

Minute numbers of contaminant CD8<sup>+</sup> T cells or CD11b<sup>+</sup>CD11c<sup>+</sup> NK cells are the source of IFN- $\gamma$  in IL-12/IL-18-stimulated mouse macrophage populations, Schleicher *et al.*, <u>Blood</u> Vol. 105, Pages 1319-1328, February 2005.

Cell Permeable Peptide of JNK Inhibitor Prevents Islet Apoptosis Immediately After Isolation and Improves Islet Graft Function, Noguchi *et al.*, American Journal of Transplantation, Vol. 5, Pages 1848-1855, August 2005.

Rescue Purification Maximizes the Use of Human Islet Preparations for Transplantation, Ichii *et al.*, <u>American Journal of Transplantation</u>, Vol. 5, Pages 21-30, © 2005.

**Prognostic Significance of Tissue Factor in Pancreatic Ductal Adenocarcinoma,** Nitori *et al.*, <u>Clinical Cancer Research</u>, Vol. 11, Pages 2531-2539, April 2005.

**Development of Colorimetric Microtiter Plate Assay for Assessment of Antimicrobials against Acanthamoeba**, McBride *et al.*, <u>American Society for Microbiology</u>, Vol. 43, Pages 629-634, September 2004.

Circulating fibroblasts, Phillips, Journal of Clinical Investigation, Vol. 114, Pages 438-446, August 2004.

Galectin-3 is induced in rheumatoid arthritis synovial fibroblasts after adhesion to cartilage oligomeric matrix protein, Neidhart *et al.*, <u>Animals of the Rheumatic Diseases</u> Pages 64:419-424, June 2004.

**Cellular engineering of ventricular adult rat cardiomyocytes**, <u>Cardiovascular Research</u>, 2003 Oct 1;59(4):874-882. "Detachment of cultured vARCs using Accutase is well compatible with ectopic gene expression and yields a viable transgenic population of vARCs that eventually may be suitable as transgenic cardiomyocyte grafts."